

**Comfortis®**  
(spinosad)

**Your veterinarian  
is the best source**  
for your pet's health information

Ask your veterinarian about your pet's healthcare and flea protection needs. They can provide the most accurate information and recommend flea control based on prevalence and seasonality in your area and the specific needs of your pet.

**Sign up for monthly  
reminders to give  
Comfortis:**

Text **COMFORTIS** to **50280**  
for texts or register for emails  
at [comfortis.com](http://comfortis.com).



**Green Chemistry**

The introduction of spinosad for use in agriculture resulted in the receipt of a Presidential Green Chemistry Challenge Award\* in 1999 from the U.S. Environmental Protection Agency. The award was presented to Dow AgroSciences LLC in the category of "Designing Safer Chemicals" for the introduction of spinosad as an insect control product for use on crops. This award demonstrates that spinosad as a technology for insect control has a favorable environmental profile. This award is not relevant to the safety and efficacy of Comfortis, nor does it confer any environmental benefit to Comfortis.



\* The Presidential Green Chemistry Challenge Awards Program, Summary of 1999 Award Entries and Recipients

Comfortis, Elanco, and the diagonal bar logo are trademarks of Eli Lilly and Company or its affiliates.

©2018 Eli Lilly and Company or its affiliates. USCACCMF00068(2)

**Elanco**

**Comfortis®**  
(spinosad)

**Fast-acting  
flea control  
for your pet**



For product label, including complete safety information, see pages 6-11.

**Elanco**

Comfortis®  
(spinosad)



## Proven protection against the threat of fleas

One tablet, once a month kills fleas and prevents and treats flea infestations in dogs and cats.\*

- **Trusted** — the first monthly oral flea control for dogs and cats
- **Fast-acting** for pets needing immediate flea relief
  - Starts killing fleas within 30 minutes<sup>1</sup>
  - In a controlled laboratory study:
    - Killed 100% of fleas in 4 hours in dogs
    - Killed 98% of fleas in 4 hours in cats
  - Proven to improve signs of flea allergy dermatitis (FAD) through its rapid action on fleas<sup>1</sup>
- **Long-lasting** 30-day protection<sup>1</sup>
  - Even if your pet swims, bathes or grooms frequently
- **Convenient** flavored tablet avoids mess
  - No need to isolate your pet from children or other pets after treatment
- **Safe** — Approved by the FDA and available by prescription only



## Important safety information

**Cats:** The most common adverse reaction recorded in clinical trials was vomiting. Other adverse reactions were lethargy, decreased appetite, weight loss, and diarrhea. Use with caution with concomitant extra-label use of ivermectin. The safe use of Comfortis in breeding, pregnant, or lactating cats has not been evaluated.

**Dogs:** The most common adverse reaction reported is vomiting. Other adverse reactions reported include lethargy, decreased appetite, incoordination, diarrhea, and seizures. Serious adverse reactions have been reported following concomitant extra-label use of ivermectin with Comfortis. Post-approval experience continues to support the safety of Comfortis when used concurrently with heartworm preventatives according to label directions. Use with caution in breeding females and dogs with pre-existing epilepsy. The safe use of Comfortis in breeding males has not been evaluated.

For product label, including complete safety information see pages 6 through 11.

\*Comfortis is approved for dogs and puppies 14 weeks and older and 5.0 lbs and greater and cats and kittens 14 weeks and older and 4.1 lbs and greater.

**Comfortis**<sup>®</sup>  
(spinosad)



## Fleas: more than a nuisance — a threat to your pet's health

Even the most conscientious pet owner can be faced with a flea infestation. A pet could be exposed in your home or backyard or when out on walks.

A few fleas can turn into a major infestation if not treated quickly. Female fleas can lay 40 to 50 eggs a day and up to 2,000 in her lifetime.<sup>2</sup> That's why it's important to kill fleas quickly, before they can start laying eggs.

- Fleas may cause extreme discomfort for your pet: scratching, chewing, biting, fur loss and restlessness.
- Severe infestations may cause anemia, particularly in young or older pets.
- Fleas are the source of flea allergy dermatitis (FAD), a common veterinary skin condition.<sup>3</sup>
- Tapeworm infections may be transmitted by fleas<sup>4</sup>



## For best flea control results

**Eliminate fleas on your pet** with a fast-acting product recommended by your veterinarian

**Eliminate fleas in your home** by vacuuming areas where your pet sits and sleeps and laundering bedding

## Prevent future infestations

Even if you don't see adult fleas, flea eggs, larvae and pupae can exist in the pet's environment and create a reinfestation.

- Monthly treatment should continue year-round<sup>5</sup>
- Treat all pets in your home with an approved flea control product
- Visit your veterinarian regularly

<sup>1</sup>Comfortis prescribing information.

<sup>2</sup>Dryden, M. 1989. Host association, on-host longevity and egg production of *Ctenocephalides felis felis*. *Vet. Parasitol.* 34:117-122.

<sup>3</sup>Sousa, C., DVM. 1997. "Fleas, flea allergy, and flea control, a review." *Dermatology Online Journal* 3.2:7. <<http://dermatology.cdlib.org>>.

<sup>4</sup><<https://www.cpcvet.org/guidelines/fleas/>>. Accessed 2/14/18.

<sup>5</sup><<https://www.cpcvet.org/guidelines/general-guidelines/>>. Accessed 2/14/18.

# Comfortis™ (spinosad)

Chewable Tablets

## COMFORTIS™ -Cats

(spinosad)

Chewable Tablets

For information on dogs, see reverse side.

**Caution:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

### Description:

COMFORTIS (spinosad) is available in three sizes of chewable flavored tablets for oral administration to cats and kittens according to their weight. Each chewable flavored tablet is formulated to provide a minimum spinosad dosage of 22.5 mg/lb (50 mg/kg). Spinosad is a member of the spinosyns class of insecticides, which are non-antibacterial tetracyclic macrolides. Spinosad contains two major factors, spinosyn A and spinosyn D, derived from the naturally occurring bacterium, *Saccharopolyspora spinosa*. Spinosyn A and spinosyn D have the chemical compositions 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[[5-(dimethylamino)tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-14-methyl-1H-as-indaceno[3,2-d]oxacyclododecin-7,15-dione and 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[[5-(dimethylamino)tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-4,14-dimethyl-1H-as-indaceno[3,2-d]oxacyclododecin-7,15-dione, respectively.

### Indications:

COMFORTIS kills fleas and is indicated for the prevention and treatment of flea infestations (*Ctenocephalides felis*), for one month, on cats and kittens 14 weeks of age and older and 4.1 pounds of body weight or greater.

### Dosage and Administration:

COMFORTIS is given orally once a month, at the minimum dosage of 22.5 mg/lb (50 mg/kg).

**Do not use the dosing schedule below when administering COMFORTIS to dogs, as it can result in an overdose.**

### Dosage Schedule:

Body Weight	Spinosad Per Tablet (mg)	Tablets Administered
4.1 to 6 lbs	140	One
6.1 to 12 lbs	270	One
12.1 to 24* lbs	560	One

\*Cats over 24 lbs should be administered the appropriate combination of tablets.

Administer COMFORTIS with food for maximum effectiveness.

COMFORTIS is a chewable tablet that can be consumed by cats when offered by the owner just prior to or after feeding. Alternatively, COMFORTIS may be offered in food or administered like other tablet medications. COMFORTIS should be administered at monthly intervals.

If vomiting occurs within an hour of administration, redose with another full dose. If a dose is missed, administer COMFORTIS with food and resume a monthly dosing schedule.

Treatment with COMFORTIS may begin at any time of the year, preferably starting one month before fleas become active and continuing monthly through the end of flea season. In areas where fleas are common,

year-round, monthly treatment with COMFORTIS should continue the entire year without interruption.

To minimize the likelihood of flea reinfestations, it is important to treat all animals within a household with an approved flea protection product.

### Contraindications:

There are no known contraindications for the use of COMFORTIS.

### Warnings:

Not for human use. Keep this and all drugs out of the reach of children.

### Precautions:

Use with caution with concomitant extra-label use of ivermectin (see ADVERSE REACTIONS).

The safe use of COMFORTIS in breeding, pregnant, or lactating cats has not been evaluated.

### Adverse Reactions:

In a well-controlled US field study, which included a total of 211 cats (139 treated with COMFORTIS and 72 treated with an active topical control once a month for 3 treatments), no serious adverse reactions were attributed to the administration of COMFORTIS.

Over the 90-day study period, all observations of potential adverse reactions were recorded. Reactions that occurred at an incidence > 1% within any of the 3 months of observations are presented in the following table. The most frequently reported adverse reaction in cats was vomiting.

Percentage of Cats (%) with Adverse Reactions

	Month 1		Month 2		Month 3	
	COMFORTIS (n=139)	Active Topical Control (n=72)	COMFORTIS (n=135)	Active Topical Control (n=69)	COMFORTIS (n=132)	Active Topical Control (n=67)
Vomiting	14.4	1.4	14.8	1.4	13.6	4.5
Lethargy	3.6	0.0	0.7	0.0	1.5	1.5
Anorexia	2.2	0.0	0.7	0.0	2.3	1.5
Weight Loss	1.4	0.0	0.0	0.0	3.0	0.0
Diarhea	1.4	1.4	0.7	2.9	2.3	1.5

Over the 3-month (3-dose) study, vomiting occurred on the day of or the day after at least one dose in 28.1% (39/139) of the cats treated with COMFORTIS and in 2.8% (2/72) of the cats treated with the active topical control. Three of the 139 cats treated with COMFORTIS vomited on the day of or the day after all three doses.

Two cats that received extra-label topical ivermectin on Day -1 of the field study developed lethargy on Day 1 after COMFORTIS administration on Day 0.

For technical assistance or to report suspected adverse drug events, contact Elanco US Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>

### Mode of Action:

The primary target of action of COMFORTIS in insects is an activation of nicotinic acetylcholine receptors (nAChRs). Spinosad does not interact with known binding sites of other nicotinic or GABAergic insecticides such as neonicotinoids, fiproles, milbemycins, avermectins, and cyclodienes. Insects treated with spinosad show involuntary muscle contractions and tremors resulting from activation of motor neurons. Prolonged spinosad-induced hyperexcitation results in prostration, paralysis, and flea death. The selective toxicity of spinosad between insects and vertebrates may be conferred by the differential sensitivity of the insect versus vertebrate nAChRs.

### Effectiveness:

In a well-controlled laboratory study, COMFORTIS began to kill fleas 30 minutes after administration and demonstrated 98% effectiveness within 4 hours. COMFORTIS kills fleas before they can lay eggs. In a separate well-controlled laboratory study, COMFORTIS demonstrated

100% effectiveness on the first day following treatment and >90% effectiveness on Day 30.

If a severe environmental infestation exists, fleas may persist for a period of time after dose administration due to the emergence of adult fleas from pupae already in the environment.

In a field study conducted in households with existing flea infestations, flea count reductions of 97.5% were observed one month after the first treatment and 99.3% after three monthly treatments with COMFORTIS. Cats with pre-existing signs of flea allergy dermatitis showed improvement in erythema, papules, scaling, alopecia, dermatitis/pyodermitis, and pruritus as a direct result of eliminating the fleas.

#### Animal Safety:

In a margin of safety study, COMFORTIS was administered orally to 14-week-old kittens at 1X, 3X, and 5X the upper half (75 – 100 mg/kg) of the therapeutic dose band for six monthly dosing intervals 28 days apart. Vomiting was observed across all groups, but was seen with greater frequency in cats in the treated groups; it did not increase with increasing doses. Loose stool was observed in all but the 3X treatment group. Food consumption was decreased in the 5X female cats. COMFORTIS was not associated with clinically significant changes in hematology, clinical chemistry, coagulation, or urinalysis parameters. Cats administered COMFORTIS once monthly for 6 months in the 3X and 5X dose groups demonstrated cytoplasmic vacuolation, consistent with phospholipidosis, in the liver, lung, and adrenal gland. The long term effects of phospholipidosis are unknown. The administration of COMFORTIS was not associated with any clinically significant, gross necropsy or histopathological changes.

In a well-controlled field study, COMFORTIS was administered safely in conjunction with other frequently used veterinary products, including tapeworm anthelmintics, antibiotics, and an approved heartworm preventative containing ivermectin. Hematology and clinical chemistry values were compared pre- and post-study and were unremarkable.

#### Storage Information:

Store at 20 to 25°C (68 to 77°F), excursions permitted between 15 to 30°C (59 to 86°F).

#### How Supplied:

COMFORTIS is available in three tablet sizes for use in cats: 140, 270 or 560 mg. Each tablet size is available in color-coded packages of 6 tablets.

## COMFORTIS™ -Dogs

(spinosad)  
Chewable Tablets

For information on cats, see reverse side.

**Caution:** Federal (USA) law restricts this drug to use by or on the order of a licensed veterinarian.

#### Description:

COMFORTIS (spinosad) is available in five sizes of chewable flavored tablets for oral administration to dogs and puppies according to their weight. Each chewable tablet is formulated to provide a minimum spinosad dosage of 13.5 mg/lb (30 mg/kg). Spinosad is a member of the spinosyns class of insecticides, which are non-antibacterial tetracyclic macrolides. Spinosad contains two major factors: spinosyn A and spinosyn D, derived from the naturally occurring bacterium, *Saccharopolyspora spinosa*. Spinosyn A and spinosyn D have the chemical compositions 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[5-dimethylamino)-tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-14-methyl-1H-as-indaceno[3,2-d]oxacyclododecin-7,15-dione and 2-[(6-deoxy-2,3,4-tri-O-methyl- $\alpha$ -L-mannopyranosyl)oxy]-13-[5-dimethylamino)-tetrahydro-6-methyl-2H-pyran-2-yl]oxy]-9-ethyl-2,3,3a,5a,5b,6,9,10,11,12,13,14,16a,16b-tetradecahydro-4,14-dimethyl-1H-as-indaceno[3,2-d]oxacyclododecin-7,15-dione, respectively.

#### Indications:

COMFORTIS kills fleas and is indicated for the prevention and treatment of flea infestations (*Ctenocephalides felis*) for one month, on dogs and puppies 14 weeks of age and older and 5.0 pounds of body weight or greater.

#### Dosage and Administration:

COMFORTIS is given orally once a month, at the recommended minimum dosage of 13.5 mg/lb (30 mg/kg).

**Do not use the dosing schedule below when administering COMFORTIS to cats, as it can result in an underdosage.**

#### Dosage Schedule:

Body Weight	Spinosad Per Tablet (mg)	Tablets Administered
5 to 10 lbs	140	One
10.1 to 20 lbs	270	One
20.1 to 40 lbs	560	One
40.1 to 60 lbs	810	One
60.1 to 120* lbs	1620	One

\* Dogs over 120 lbs should be administered the appropriate combination of tablets. Administer COMFORTIS with food for maximum effectiveness.

COMFORTIS is a chewable tablet and is readily consumed by dogs when offered by the owner just prior to feeding. Alternatively, COMFORTIS may be offered in food or administered like other tablet medications. COMFORTIS should be administered at monthly intervals.

If vomiting occurs within an hour of administration, redose with another full dose. If a dose is missed, administer COMFORTIS with food and resume a monthly dosing schedule.

Treatment with COMFORTIS may begin at any time of the year, preferably starting one month before fleas become active and continuing monthly through the end of flea season. In areas where fleas are common year-round, monthly treatment with COMFORTIS should continue the entire year without interruption.

To minimize the likelihood of flea reinfestation, it is important to treat all animals within a household with an approved flea protection product.

#### Contraindications:

There are no known contraindications for the use of COMFORTIS.

#### Warnings:

Not for human use. Keep this and all drugs out of the reach of children.

Serious adverse reactions have been reported following concomitant extra-label use of ivermectin with COMFORTIS (see POST APPROVAL EXPERIENCE).

#### Precautions:

COMFORTIS is for use in dogs and puppies 14 weeks of age and older (see ANIMAL SAFETY).

Use with caution in breeding females (see ANIMAL SAFETY). Use with caution in dogs with pre-existing epilepsy (see ADVERSE REACTIONS). The safe use of COMFORTIS in breeding males has not been evaluated.

#### Adverse Reactions:

In a well-controlled US field study, which included a total of 470 dogs (330 dogs treated with COMFORTIS and 140 dogs treated with an active control), no serious

adverse reactions were observed with COMFORTIS. All reactions were regarded as mild and did not result in any dog being removed from the study.

Over the 90-day study period, all observations of potential adverse reactions were recorded. Reactions that occurred at an incidence > 1% within any of the 3 months of observation are presented in the following table. The most frequently reported adverse reaction in dogs in the COMFORTIS and active control groups was vomiting. The occurrence of vomiting, most commonly within 48 hours after treatment, decreased with repeated doses of COMFORTIS.

Percentage of Dogs (%) with Adverse Reactions

	Month 1		Month 2		Month 3	
	COMFORTIS Chewable Tablets (N=330)	Active Topical Control (N=139) <sup>a</sup>	COMFORTIS Chewable Tablets (N=282)	Active Topical Control (N=124)	COMFORTIS Chewable Tablets (N=260)	Active Topical Control (N=125)
Vomiting	12.7	12.2	7.8	3.2	5.8	4.8
Decreased Appetite	9.1	5.0	2.8	1.6	1.9	0.8
Lethargy	7.6	5.0	3.5	4.0	1.2	0.8
Diarrhea	6.7	5.0	4.3	0.8	1.2	0.0
Cough	3.9	5.0	0.4	2.4	0.0	0.0
Polydipsia	2.4	1.4	0.7	0.0	0.4	0.0
Vocalization	1.8	0.0	0.4	0.0	0.4	0.0
Increased Appetite	1.5	0.0	0.4	0.8	0.4	0.0
Erythema	1.5	0.0	0.4	0.0	0.4	0.0
Hyperactivity	1.2	1.4	0.0	0.0	0.4	0.0
Excessive Salivation	1.2	0.0	0.4	0.0	0.0	0.0

<sup>a</sup> This number (n=139) is less than the total number of dogs in the safety population for the active control group (n=140) because one dog joined the study late and was only dosed at Month 3.

In US and European field studies, no dogs experienced seizures when dosed with COMFORTIS at the therapeutic dose range of 13.5-27.3 mg/lb (30-60 mg/kg), including 4 dogs with pre-existing epilepsy. Four epileptic dogs that received higher than the maximum recommended dose of 27.3 mg/lb (60 mg/kg) experienced at least one seizure within the week following the second dose of COMFORTIS, but no seizures following the first and third doses. The cause of the seizures observed in the field studies could not be determined.

#### Post Approval Experience (June 2009)

The following adverse reactions are based on post-approval adverse drug event reporting. The adverse reactions are listed in decreasing order of frequency: vomiting, depression/lethargy, anorexia, ataxia, diarrhea, pruritus, trembling, hypersalivation and seizures.

Following concomitant extra-label use of ivermectin with COMFORTIS, some dogs have experienced the following clinical signs: trembling/twitching, salivation/drooling, seizures, ataxia, mydriasis, blindness and disorientation.

Post approval experience continues to support the safety of COMFORTIS when used concurrently with heartworm preventatives according to label directions.

For technical assistance or to report suspected adverse drug events, contact Elanco US Inc. at 1-888-545-5973. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/AnimalVeterinary/SafetyHealth>

#### Mode of Action:

The primary target of action of COMFORTIS in insects is an activation of nicotinic acetylcholine receptors (nAChRs). Spinosad does not interact with known insecticidal binding sites of other nicotinic or GABAergic insecticides such as neonicotinoids, fiproles, milbemycins, avermectins, and cyclodienes. Insects treated with spinosad show involuntary muscle contractions and tremors resulting from activation of motor neurons. Prolonged spinosad-induced hyperexcitation results in prostration, paralysis, and flea death. The selective toxicity of spinosad between insects and vertebrates may be conferred by the differential sensitivity of the insect versus vertebrate nAChRs.

#### Effectiveness:

In a well-controlled laboratory study, COMFORTIS began to kill fleas 30 minutes after administration and demonstrated 100% effectiveness within 4 hours. COMFORTIS kill fleas before they can lay eggs. If a severe environmental infestation exists, fleas may persist for a period of time after dose administration due to the emergence of adult fleas from pupae already in the environment. In field studies conducted in households with existing flea infestations of varying severity, flea reductions of 98.0% to 99.8% were observed over the course of 3 monthly treatments with COMFORTIS. Dogs with signs of flea allergy dermatitis showed improvement in erythema, papules, scaling, alopecia, dermatitis/pyodermitis and pruritus as a direct result of eliminating the fleas.

#### Animal Safety:

COMFORTIS was tested in pure and mixed breeds of healthy dogs in well-controlled clinical and laboratory studies. No dogs were withdrawn from the field studies due to treatment-related adverse reactions.

In a dose tolerance study, COMFORTIS was administered orally to adult Beagle dogs at average doses of up to 100 mg/kg once daily for 10 consecutive days (16.7 times the maximum recommended monthly dose). Vomiting was seen in 5 of 6 treated dogs during the first 6 days of treatment, usually within 2.5 hours of dosing. Treated females lost weight early in the treatment period, but their weights were similar to control dogs by the end of the 24-day study. COMFORTIS was not associated with any clinically significant changes in hematology, blood coagulation or urinalysis parameters; however, mild elevations in ALT occurred in all dogs treated with COMFORTIS. By day 24, ALT values had returned to near baseline levels. Phospholipidosis (vacuolation) of the lymphoid tissue was seen in all dogs treated with COMFORTIS, the long-term effects of which are unknown.

In a margin of safety study, COMFORTIS was administered orally to 6-week-old Beagle puppies at average doses of 1.5, 4.4, and 7.4 times the maximum recommended dose at 28-day intervals over a 6-month period. Vomiting was observed across all groups, including the control. Increased vomiting was observed at elevated doses, usually within 1 hour following administration. Vomiting at all doses decreased over time and stabilized when puppies were 14 weeks of age. The average daily and total weight gains of treated dogs were smaller than control dogs and were dose dependent. COMFORTIS was not associated with clinically significant changes in hematology, clinical chemistry, coagulation or urinalysis parameters. Phospholipidosis (vacuolation) of the lymphoid tissue was seen in some dogs in the 4.4X group and all dogs in the 7.4X group. The long term effects of phospholipidosis are unknown. Treatment with COMFORTIS was not associated with any other clinically significant adverse clinical observations, gross necropsy or histopathological changes.

In a reproductive safety study, COMFORTIS was administered orally to female Beagles at 1.3 and 4.4 times the maximum recommended therapeutic dose every 28 days prior to mating, during gestation, and during a six-week lactation period. No treatment-related adverse effects were noted for conception rates in the dams, or for mortality, body temperature, necropsy, or histopathology findings for the dams or puppies. One dam from each treatment group experienced early pregnancy loss and one additional high dose dam aborted late term. The treated dams experienced more vomiting, especially at one hour post-dose, than the control dams. Puppies from dams treated at 1.3 times the maximum recommended therapeutic dose had lower body weights than puppies from control dams. Although puppy mortality between treated and control dams was not different, the puppies from the treated dams experienced more lethargy (4.4X group only), dehydration, weakness and felt cold to the touch (4.4X group only) than puppies from control dams.

A pilot study without a control group was conducted to analyze milk from three lactating dogs treated with an experimental formulation of spinosad at 1.5 times the maximum recommended dose administered at day 28 of gestation and 24 hours prior to parturition. The data demonstrated that spinosad was excreted in the milk of these dogs. Mortality and morbidity were greatest in puppies from the dam with the highest spinosad level in milk. The spinosad milk: reference plasma exposure ratio calculated from this study ranged from 2.2 to 3.5.

In well-controlled field studies, COMFORTIS was administered safely in conjunction with other frequently used veterinary products, such as vaccines, anthelmintics, antibiotics, steroids, flea and tick control products, anesthetics, NSAIDs, antihistamines, alternative/herbal remedies, shampoos, and prescription diets. Changes in hematology, clinical chemistry and urinalysis values were compared pre- and post-study and were unremarkable.

#### Storage Information:

Store at 20-25°C (68-77°F), excursions permitted between 15 to 30°C (59 to 86°F).

#### How Supplied:

COMFORTIS is available in five tablet sizes for use in dogs: 140, 270, 560, 810 or 1620 mg. Each tablet size is available in color-coded packages of 6 tablets.

NADA #141-277, Approved by the FDA

Manufactured for:

Elanco US Inc.  
Greenfield, IN 46140  
[www.comfortis.com](http://www.comfortis.com)

Elanco, Comfortis and the diagonal bar are trademarks owned or licensed by

El Lilly and Company, its subsidiaries or affiliates.

**Elanco**

NDC 58198-4222-6  
NDC 58198-4223-6  
NDC 58198-4224-6  
NDC 58198-4225-6  
NDC 58198-4227-6

CA4222  
CA4223  
CA4224  
CA4225  
CA4227

03B431  
JUN 2017  
CMF1a